arXiv:1210.1216v3 [math.NT] 15 Oct 2012

EULER PRODUCTS BEYOND THE BOUNDARY

TARO KIMURA*, SHIN-YA KOYAMA, AND NOBUSHIGE KUROKAWA

Abstract. For the Riemann zeta and the Dirichlet L-functions, we investigate their behavior of the Euler products on the critical line. A refined version of the Riemann hypothesis, which is named "the Deep Riemann Hypothesis" (DRH), is proposed. We prove that the analogue of the DRH is true for the function field cases.

1. Introduction

Let χ be a primitive Dirichlet character with conductor N. The Dirichlet L-function is expressed by the Euler product

$$L(s,\chi) = \prod_p (1-\chi(p)p^{-s})^{-1}, \tag{1}$$

where p runs through all primes. The product (1) is absolutely convergent in $\mathrm{Re}(s) > 1$. It is known that $L(s,\chi)$ has a meromorphic continuation to all $s \in \mathbf{C}$, which is entire if $\chi \neq \mathbf{1}$, and has a simple pole at $s = 1$ if $\chi = \mathbf{1}$.

In this paper we study the values $L(s,\chi)$ beyond the boundary $\mathrm{Re}(s) = 1$ of the absolute convergence region $\mathrm{Re}(s) > 1$ from the viewpoint of its relation to the values of the Euler product. Few results are known along this context. The classical results concerning the fact that the Euler product (1) converges to $L(1+it,\chi)$ ($t \in \mathbf{R},\ t \neq 0$) are seen in the textbooks for either case $\chi = \mathbf{1}$ ([T] Chapter 3) or $\chi \neq \mathbf{1}$ ([M]). The only work we can find beyond these is that of Goldfeld [G], Kuo-Murty [KM] and Conrad [C]. Goldfeld [G] and Kuo-Murty [KM] dealt with the L-functions of elliptic curves at $s = 1$ with their results supporting the Birch and Swinnerton-Dyer conjecture. Conrad [C] treated more general Euler products for $\mathrm{Re}(s) \geq 1/2$.

The (generalized) Riemann Hypothesis (GRH) for $L(s,\chi)$ asserts that $L(s,\chi) \neq 0$ in $\mathrm{Re}(s) > 1/2$. When $\chi \neq \mathbf{1}$, it is equivalent to the following conjecture.

Conjecture 1. *If* $\chi \neq \mathbf{1}$, *then for* $\mathrm{Re}(s) > 1/2$ *we have*

$$L(s,\chi) = \lim_{n\to\infty} \prod_{p \leq n} (1-\chi(p)p^{-s})^{-1},$$

where the product is taken over all primes p *satisfying* $p \leq n$.

Note that the order of primes which participate in the product is important, because it is not absolutely convergent.

Here we examine a "deeper" conjecture in the sense that we dig into the line $\mathrm{Re}(s) = 1/2$.

2000 *Mathematics Subject Classification.* 11M06.

Key words and phrases. The Riemann zeta function; Dirichlet L-functions; the Riemann hypothesis; the generalized Riemann hypothesis; Euler products.

*Partially supported by JSPS Research Fellowships for Young Scientists (No. 23-593).

Conjecture 2 (Deep Riemann Hypothesis (DRH))**.** *If* $\chi \neq \mathbf{1}$ *and* $L(s,\chi) \neq 0$ *with* $\mathrm{Re}(s) = \frac{1}{2}$*, we have*

$$\lim_{n\to\infty} \prod_{p\leq n} (1-\chi(p)p^{-s})^{-1} = L(s,\chi) \times \begin{cases} \sqrt{2} & (s=\frac{1}{2} \text{ and } \chi^2 = \mathbf{1}) \\ 1 & (\text{otherwise}) \end{cases},$$

where the product is taken over all primes p satisfying $p \leq n$.

The prototype version of this conjecture was proposed in [C]. For a generalization of Conjecture 2 to the case including $\chi = \mathbf{1}$, see Akatsuka [A].

It is seen that Conjecture 2 implies Conjecture 1, but that the converse is not true. It is an easy task to obtain numerical support of Conjecture 2, since the convergence of the left hand side is fairly fast.

2. Function Field Analogs

In this section, we prove an analog of Conjecture 2 for function fields of one variable over a finite field.

Let $\mathbf{F}_p$ be the finite field of p elements. We fix a conductor $f(T) \in \mathbf{F}_p[T]$ and introduce a "Dirichlet" character

$$\chi : \ (\mathbf{F}_p[T]/(f))^\times \to \mathbf{C}^\times.$$

We define the "Dirichlet" L-function by the Euler product:

$$L_{\mathbf{F}_p(T)}(s,\chi) = \prod_h (1-\chi(h)N(h)^{-s})^{-1},$$

where $h = h(T) \in \mathbf{F}_p[T]$ runs through monic irreducible polynomials, and $N(h) = p^{\deg h}$. By the celebrated work of Kornblum [K], it is proved that the above Euler product is absolutely convergent in $\mathrm{Re}(s) > 1$, and is a polynomial in p^{-s} of degree less than $\deg f$ if $\chi \neq \mathbf{1}$.

We prove the following theorem.

Theorem 1 (DRH over function fields)**.** *Let p, f and χ be as above. Put $K = \mathbf{F}_p(T)$ and assume $\chi \neq \mathbf{1}$. Then the following (1) and (2) are true.*

(1) *For* $\mathrm{Re}(s) > 1/2$*, we have*

$$\lim_{n\to\infty} \prod_{\deg h \leq n} (1-\chi(h)N(h)^{-s})^{-1} = L_K(s,\chi).$$

(2) *For* $t \in \mathbf{R}$ *with* $L_K(\frac{1}{2}+it,\chi) \neq 0$*, it holds that*

$$\lim_{n\to\infty} \prod_{\deg h \leq n} (1-\chi(h)N(h)^{-\frac{1}{2}-it})^{-1} = L_K\left(\frac{1}{2}+it,\chi\right) \times \begin{cases} \sqrt{2} & (\chi^2 = \mathbf{1},\ t \in \frac{\pi}{\log p}\mathbf{Z}) \\ 1 & (\text{otherwise}) \end{cases}.$$

Proof. It suffices to prove (2), since (2) implies (1). We estimate the product

$$E_n = \prod_{\deg h \leq n} \left(1-\chi(h)N(h)^{-\frac{1}{2}-it}\right)^{-1}$$

by dealing with its logarithm

$$\log E_n = \sum_{\deg h \leq n} \sum_{k=1}^{\infty} \frac{\chi(h)^k}{k} p^{-k(\frac{1}{2}+it)\deg h}.$$

We divide the sum into three parts as

$$\log E_n = A(n) + B(n) + C(n)$$

with

$$\begin{aligned} A(n) &= \sum_{k=1}^{\infty} \sum_{\deg h \le n/k} \frac{\chi(h)^k}{k} p^{-k(\frac{1}{2}+it)\deg h}, \\ B(n) &= \sum_{n/2 \le \deg h \le n} \frac{\chi(h)^2}{2} p^{-2(\frac{1}{2}+it)\deg h}, \\ C(n) &= \sum_{k=3}^{\infty} \sum_{n/k < \deg h \le n} \frac{\chi(h)^k}{k} p^{-k(\frac{1}{2}+it)\deg h}. \end{aligned}$$

By the above mentioned Kornblum's theorem, we put

$$L_K(s,\chi) = \prod_{j=1}^{r} (1 - \lambda_j p^{-s})$$

with $|\lambda_j| = \sqrt{p}$ or 1. Then by taking the logarithmic derivatives of

$$\prod_h (1 - \chi(h) N(h)^{-s})^{-1} = \prod_{j=1}^{r} (1 - \lambda_j p^{-s}) \quad (\mathrm{Re}(s) > 1)$$

and comparing the coefficients of p^{-sk}, we have

$$\sum_{(\deg h)|k} (\deg h) \chi(h)^{\frac{k}{\deg h}} = -\sum_{j=1}^{r} \lambda_j^k \quad (k \ge 1).$$

By this identity, the first partial sum $A(n)$ is calculated as

$$\begin{aligned} A(n) &= \sum_{k \le n} \frac{p^{-(\frac{1}{2}+it)k}}{k} \sum_{(\deg h)|k} (\deg h) \chi(h)^{\frac{k}{\deg h}} \\ &= -\sum_{j=1}^{r} \sum_{k=1}^{n} \frac{1}{k} \left(\frac{\lambda_j}{p^{\frac{1}{2}+it}} \right)^k. \end{aligned}$$

By the Kornblum theorem we have $\left| \frac{\lambda_j}{p^{\frac{1}{2}+it}} \right| \le 1$ and the assumption $L_K(\frac{1}{2}+it,\chi) \ne 0$ tells that $\frac{\lambda_j}{p^{\frac{1}{2}+it}} \ne 1$. Hence it holds that

$$\begin{aligned} \lim_{n\to\infty} A(n) &= \sum_{j=1}^{r} \log \left(1 - \frac{\lambda_j}{p^{\frac{1}{2}+it}} \right) \\ &= \log L_K \left(\frac{1}{2} + it, \chi \right). \end{aligned}$$

Next for estimating $B(n)$, we use the well-known fact that

$$\sum_{\deg h < n} \frac{1}{N(h)} \sim \log n \qquad (n \to \infty).$$

When $\chi^2 = \mathbf{1}$ and $t \in \frac{\pi}{\log p}\mathbf{Z}$, we compute that

$$\begin{aligned} B(n) &= \frac{1}{2} \sum_{n/2 \le \deg h \le n} p^{-(1+2it)\deg h} \\ &= \frac{1}{2}\left(\sum_{1 \le \deg h \le n} p^{-(1+2it)\deg h} - \sum_{1 \le \deg h < n/2} p^{-(1+2it)\deg h} \right) \\ &= \frac{1}{2}\left(\left(\log n + C + O(n^{-1})\right) - \left(\log \frac{n}{2} + C + O(n^{-1})\right)\right) \\ &= \frac{1}{2}\left(\log 2 + O(n^{-1})\right). \end{aligned}$$

Hence

$$\lim_{n \to \infty} B(n) = \log \sqrt{2}.$$

Finally, $C(n) \to 0$ as $n \to \infty$, because $\sum_h p^{-s \deg h}$ is absolutely convergent in $\mathrm{Re}(s) > 1$. Now that $k \geq 3$, we have $|p^{-k(\frac{1}{2}+it)\deg h}| \leq p^{\frac{3}{2}}$. ☐

Conjecture 2 and Theorem 1 are generalized to automorphic L-functions. See Lownes [L].

3. Numerical Calculations

In this section we show some numerical datum which are admitted as evidence of the Deep Riemann Hypothesis (Conjecture 2). According to Conjecture 2, even if we consider along the critical line $\mathrm{Re}(s) = 1/2$, the Euler product gives a meaningful value such as $L(s, \chi)$ or $\sqrt{2}L(s, \chi)$. If this is true, the partial Euler product

$$L_x(s, \chi) = \prod_{p \leq x} (1 - \chi(p)p^{-s})^{-1},$$

converges to $L(s, \chi)$ or $\sqrt{2}L(s, \chi)$ as $x \to \infty$ even on the critical line $\mathrm{Re}(s) = 1/2$. We formally put $L_x(s, \chi) = L(s, \chi)$ for $x = \infty$.

First we give Table 1, which shows the accuracy of Conjecture 2 at $s = 1/2$. We find that the ratio of $\sqrt{2}L(\frac{1}{2}, \chi)$ and $L_x(\frac{1}{2}, \chi)$ is almost equal to 1 for $x = 10^7$, when χ is quadratic.

In what follows we put χ_{7a} and χ_{7b} to be the character χ modulo 7 with $\chi^2 \neq \mathbf{1}$ and $\chi^2 = \mathbf{1}$, respectively. Namely, if we define the character χ modulo 7 by giving the value at the primitive root $3 \in \mathbf{Z}/7\mathbf{Z}$, we define $\chi_{7a}(3) = \exp(\pi\sqrt{-1}/3)$ and $\chi_{7b}(3) = -1$. We also denote by χ_3 the nontrivial character modulo 3, which satisfies $\chi_3^2 = 1$.



Figure 1. Real part (left) and imaginary part (right) of $L_x(1/2 + it, \chi_{7a})$

d	$\sqrt{2}L$	E	$(\sqrt{2}L)/E$
-3	0.680049	0.688002	0.988440
-4	0.944258	0.945909	0.998254
5	0.327745	0.320619	1.022223
-7	1.621517	1.640320	0.988536
8	0.528479	0.539992	0.978680
-8	1.556230	1.521663	1.022716
-11	1.402301	1.342967	1.044181
12	0.705066	0.729170	0.966942
13	0.621678	0.618558	1.005044
-15	2.612093	2.791265	0.935809
17	1.020601	1.066235	0.957201
-19	1.137621	1.173052	0.969795
-20	2.375413	2.356696	1.007942
21	0.703235	0.724051	0.971250
-23	3.472406	3.320551	1.045732
24	1.003325	1.057376	0.948881
-24	2.223023	2.130498	1.043428
28	1.162994	1.199957	0.969196
29	0.658655	0.683281	0.963958

TABLE 1. $L := L\left(\frac{1}{2}, \left(\frac{d}{\cdot}\right)\right)$, $E := \displaystyle\prod_{p \le 10^7} \left(1 - \left(\frac{d}{p}\right)\frac{1}{\sqrt{p}}\right)^{-1}$.



FIGURE 2. Real part (left) and imaginary part (right) of $L_x(1/2 + it, \chi_{7b})$



FIGURE 3. Real part (left) and imaginary part (right) of $L_x(3/4 + it, \chi_{7a})$

Denote by p_n the n-th prime number. Figures 1, 2, 3, 4, 5 and 6 show the datum for the values

$$L_x\left(\frac{1}{2} + it,\, \chi\right), \qquad L_x\left(\frac{3}{4} + it,\, \chi\right), \qquad L_x\left(1 + it,\, \chi\right)$$



FIGURE 4. Real part (left) and imaginary part (right) of $L_x(3/4+it,\chi_{7b})$



FIGURE 5. Real part (left) and imaginary part (right) of $L_x(1+it,\chi_{7a})$



FIGURE 6. Real part (left) and imaginary part (right) of $L_x(1+it,\chi_{7b})$

for $x = p_{10}$ (green), $x = p_{100}$ (blue), $x = p_{1000}$ (yellow) and ∞ (red). Figures 1, 3 and 5 are for χ_{7a}, and Figures 2, 4 and 6 for χ_{7b}. This shows as $t \to 0$, we apparently see $L_x\left(1/2+it,\,\chi\right) \to L(1/2,\chi)$ for $\chi^2 \neq \mathbf{1}$, $L_x\left(1/2+it,\,\chi\right) \to \sqrt{2}L(1/2,\chi)$ for $\chi^2 = \mathbf{1}$, and $L_x\left(3/4+it,\,\chi\right) \to L(3/4,\chi)$, $L_x\left(1+it,\,\chi\right) \to L(1,\chi)$ for $\chi^2 = \mathbf{1}$ and $\chi^2 \neq \mathbf{1}$. This supports the DRH (Conjecture 2).

We introduce the following error function in order to estimate the speed of convergence for $L_x(s,\chi)$,

$$\delta L_x(s,\chi) = \begin{cases} \left|\frac{L_x(s,\chi)-\sqrt{2}L(s,\chi)}{\sqrt{2}L(s,\chi)}\right| & (s = 1/2 \ \mathit{and}\ \chi^2 = \mathbf{1}) \\ \left|\frac{L_x(s,\chi)-L(s,\chi)}{L(s,\chi)}\right| & (\mathit{otherwise}) \end{cases}$$

Figure 7 shows the values of $\delta L_x(s,\chi)$. When we approximate the error function as $\delta L_x(s,\chi) \sim x^{-\alpha}$, the exponents are determined by fitting the numerical results (Table 2). We see the speed of convergence becomes faster as the real part of argument gets larger.

s	$\alpha\ (\chi_{7a})$	$\alpha\ (\chi_{7b})$
1/2	0.1167	0.1978
3/4	0.3814	0.3106
1	0.6389	0.6302

TABLE 2. Exponents of $\delta L_x(s,\chi) \sim x^{-\alpha}$ for χ_{7a} and χ_{7b}.



FIGURE 7. $\delta L_x(s,\chi)$ for $s = 1/2$ (red), $s = 3/4$ (green) and $s = 1$ (blue) with χ_{7a} (left) and χ_{7b} (right)

In Figures 8, 9 and 10, the blue curves show the values

$$\rho_x(t) = \frac{1}{\pi}\mathrm{Im}\frac{d}{dt}\log L_x\left(\frac{1}{2}+it,\ \chi\right) \tag{2}$$

with $x = p_{1000}$ for χ_3, χ_{7a}, χ_{7b}, respectively. The red curves are $|L\left(\frac{1}{2}+it,\ \chi\right)|$. This function is an analog of the eigenvalue density function in random matrix theory. ζ-function on the critical line $s = 1/2 + it$ can be seen as a characteristic polynomial of a certain infinite dimensional matrix [KS, BH]

$$e^{i\vartheta(t)}\zeta\left(\frac{1}{2}+it\right) = \prod_{j=1}^{\infty}(t-t_j),$$

where $\vartheta(t)$ is the Riemann-Siegel theta function

$$\vartheta(t) = \mathrm{Im}\log\Gamma\left(\frac{it}{2}-\frac{3}{4}\right) - \frac{t}{2}\log\pi.$$

There is also a similar representation for L-function on the critical line. Thus when we define the density function of the nontribial zeros on the critical line as

$$\begin{aligned}\rho(t) &= \sum_{j=1}^{\infty}\delta(t-t_j)\\ &= \frac{1}{\pi}\mathrm{Im}\sum_{j=1}^{\infty}\frac{1}{x-x_j}\\ &= \frac{1}{\pi}\mathrm{Im}\frac{d}{dt}\log\prod_{j=1}^{\infty}(t-t_j),\end{aligned}$$

the function (2) should converge to this density function in the limit of $x \to \infty$, up to the factor coming form $\vartheta(t)$. Apparently the location of the zeros of $|L(\frac{1}{2}+it,\ \chi)|$ agrees to that of the peaks of $\rho_x(t)$ in Figures 8, 9 and 10. This suggests that the first few primes already know the nontrivial zeros of $L(s,\ \chi)$, and that the Euler product is meaningful beyond the boundary. The leaps in the red curves correspond to the zeros of $L(s,\chi)$. We normalize that the average spacings of zeros are equal to one. This reflects the fact that the multiplicity of such zeros are all one. In other words, if we express their derivatives by the Dirac's delta function, the coeffients are one. We also observe that the blue curve oscillates if and only if $\chi^2 = \mathbf{1}$. The behavior near $t = 0$ depends on whether $\chi^2 = \mathbf{1}$ or not.



FIGURE 8. $\rho_x(t)$ for χ_3



FIGURE 9. $\rho_x(t)$ for χ_{7a}

FIGURE 10. $\rho_x(t)$ for χ_{7b}

Figures 11, 12 and 13 indicate the values

$$R_x(t) = \frac{1}{\pi}\mathrm{Im}\log L_x\left(\frac{1}{2} + it,\, \chi\right)$$

for χ_3, χ_{7a}, χ_{7b}, respectively, for $x = p_{10}$ (green), $x = p_{100}$ (blue), $x = p_{1000}$ (yellow) and ∞ (red). This also seems to reflect the property of DRH. The green, blue and yellow curves appear to converge to the red one more smoothly only when $\chi^2 \neq \mathbf{1}$ (Figure 12). In the other two cases, the curves oscillate many times near the origin.

These figures also tell us that the values $\mathrm{Im}\log L(\frac{1}{2} + it)$ are almost stable for nontrivial zeros $\frac{1}{2} + it$ of the L-function, no matter how many prime numbers we take into account. This suggests that the nontrivial zeros are analogs of the critical points in physics.

Finally, Figure 14 show how the peaks of $\rho(t)$ with the smallest t in Figures 8, 9 and 10 get closer to the zeros of $L(s, \chi)$ under the same settings in Figures 11, 12 and 13.

REFERENCES

[A] H. Akatsuka: The Euler product for the Riemann zeta-function on the critical line. (preprint, 2012)
[BH] E. Brézin and S. Hikami: Characteristic polynomials of random matrices, Comm. Math. Phys. **214** (2000) 111–135.
[C] K. Conrad: Partial Euler products on the critical line, Canad. J. Math. **57** (2005) 267–297.
[G] D. Goldfeld: Sur les produits partiels eulériens attachés aux courbes elliptiques, C. R. Acad. Sci. Paris Sér. I Math. **294** (1982) 471–474.
[K] H. Kornblum: Über die Primfunktionen in einer arithmetischen Progression, Math. Zeit., **5**, 100–111 (1919).



FIGURE 11. $R_x(t)$ for χ_3



FIGURE 12. $R_x(t)$ for χ_{7a} FIGURE 13. $R_x(t)$ for χ_{7b}



FIGURE 14. Peaks in $\rho(t)$ with the smallest t for χ_3 (left), χ_{7a} (center) and χ_{7b} (right)

[KM] W. Kuo and R. Murty: On a conjecture of Birch and Swinnerton-Dyer, Canad. J. Math. **57** (2005) 328-337.

[KS] J. P. Keating and N. C. Snaith: Random matrix theory and $\zeta(1/2+it)$, Comm. Math. Phys. **214** (2000) 57–89.

[L] C. Lownes: Deep Riemann hypothesis for $GL(n)$. (preprint, 2012)

[M] F. Mertens: Ein Beitrag zur analytischen Zahlentheorie, J. Reine Angew. Math. **78** (1874) 46–72.

[T] E. C. Titchmarsh: The theory of the Riemann zeta function, Oxford University Press, 1987.

MATHEMATICAL PHYSICS LABORATORY, RIKEN NISHINA CENTER, 2-1 HIROSAWA, WAKO, SAITAMA, 351-0198, JAPAN.

E-mail address: `taro.kimura@riken.jp`

Department of Biomedical Engineering, Toyo University, 2100 Kujirai, Kawagoe, Saitama, 350-8585, Japan.

E-mail address: `koyama@toyo.jp`

Department of Mathematics, Tokyo Institute of Technology, 2-12-1 Oh-okayama, Meguro-ku, Tokyo 152-8551, Japan.

E-mail address: `kurokawa@math.titech.ac.jp`